

Application Of Modern Wound Care Using Moist Wound Healing Techniques In Diabetes Mellitus Patients

by Rafika Nur Siregar

Submission date: 03-May-2024 02:50PM (UTC+0700)

Submission ID: 2369678690

File name: Moist_Wound_Healing_Techniques_In_Diabetes_Mellitus_Patients.pdf (348.77K)

Word count: 2796

Character count: 15062



Application Of Modern Wound Care Using Moist Wound Healing Techniques In Diabetes Mellitus Patients

Rafika Nur Siregar

Nursing Diploma Study Program, Sekolah Tinggi Ilmu Kesehatan Malahayati Medan

Correspondence email: rafikanur47@gmail.com

Abstract. Diabetes Mellitus often becomes a serious health problem in various developing and developed countries in the world. Most people with diabetes mellitus are at risk of developing diabetic ulcers. If not treated quickly it can develop into an infection. The newest wound healing methods currently have an important role in treating diabetic ulcers, one of which is Modern Dressing with Moist Wound Healing. The aim of this research is to identify the effectiveness of modern wound care using moist wound healing dressing techniques in healing diabetic ulcers at the Sering Road. The research design used in this study used a quasi-experimental design with a cross-sectional method approach. The sampling used was accidental sampling, namely 30 respondents. Based on the results of the Paired T-Test and Independent T-Test, the p-value was obtained ($0.000 < \alpha 0.05$), in the control group, p value $\alpha = 0.080$ ($p > 0.05$). Thus, it can be said that there is no effectiveness of modern wound care using moist wound healing dressing techniques in healing diabetic ulcers at the Sering Road. It is hoped that Sering Road can improve the quality and quality of modern wound care services using moist wound healing dressing techniques for diabetic ulcer sufferers.

Keywords: Diabetic Ulcers, Modern Wound Care, Moist Wound Healing

INTRODUCTION

Diabetes Mellitus is a metabolic disorder characterized by an increase in blood glucose due to a decrease in insulin secretion by pancreatic beta cells (Siregar, H.K., Butar Butar, S., Pangaribuan, S.M., Siregar, S.W., Batubara, 2023). The main risks commonly found in every diabetes mellitus sufferer include hypoglycemia, hyperglycemia, diabetic ketoacidosis, dehydration and thrombosis (Rusdi, 2020).

The Diabetes Mellitus criteria in the 2018 Riskesdas refer to the consensus of the Indonesian Endocrinology Association (PERKENI) which adopted the American Diabetes Association (ADA) criteria. According to these criteria, Diabetes Mellitus is confirmed if the fasting blood glucose level is ≥ 126 mg/dl, or blood glucose 2 hours after loading ≥ 200 mg/dl, or instant blood glucose ≥ 200 mg/dl with symptoms of frequent hunger, frequent thirst, frequent urination, small amounts of water, and weight loss (Ministry of Health of the Republic of Indonesia, 2018).

As the healing time for diabetes and microangiopathy takes longer, diabetic neuropathy can cause ulcers on the feet, deformity and can result in amputation. Foot ulcers in neuropathy often occur on the plantar surface of the foot, namely in areas that experience high pressure, such as areas covering the metatarsal heads or other areas covering bone deformities. Diabetic foot ulcers contribute to $> 50\%$ of diabetic foot ulcers and often do not cause pain accompanied by bruising. One of the complications of Diabetes Mellitus is neuropathy.

Received March 30, 2024; Accepted May 03, 2024; Published June 31, 2024

* Rafika Nur Siregar, rafikanur47@gmail.com

Reduced sensation in the feet and is often associated with foot injuries. Peripheral neuropathy can cause loss of sensation in the distal areas of the feet which has a high risk of developing foot ulcers and can even lead to amputation (Siregar & Siregar, 2022).

Based on data from the International Diabetes Federation (IDF), which states that since 2020, countries in the Arab-North African region and the Western Pacific have ranked first and second with the highest prevalence of diabetes in the population aged 20-79 years among 7 regions in the world, namely of 12.2% and 11.4%. The Southeast Asia region where Indonesia is located is ranked third with a prevalence of 11.3%. Indonesia is the only country in Southeast Asia on the list, so it can be estimated that Indonesia's contribution to the prevalence of diabetes cases in Southeast Asia (Indonesian Ministry of Health, 2020).

Based on data from the International Diabetes Foundation (IDF), it is stated that there were 1785 cases of Diabetes Mellitus sufferers who experienced complications of neuropathy (63.5%), retinopathy (42%), nephropathy (7.3%), macrovascular (16%), microvascular (6%), diabetic foot wounds (15%) (Siregar et al., 2021). The majority of respondents carried out sterile wound care (53.3%), respondents carried out wound care with tools and materials that were less available or incomplete (76.7%), and the majority of respondents carried out wound care not appropriate for diabetic wound care procedures (73.3%). While the prevalence of diabetic ulcer sufferers in the United States is 15-20%, the risk of amputation is 15-46 times higher compared to sufferers who do not experience diabetes mellitus.

Diabetic ulcer sufferers in the United States require high costs for treatment each year for Diabetes Mellitus sufferers. The prevalence of diabetic ulcer sufferers in Indonesia is around 15%, the amputation rate is 30%, the mortality rate is 32%, and diabetic ulcers are the most common cause of hospital admission at 80% for Diabetes Mellitus sufferers (Rassi, Y.E.T., Efendy, 2023). Based on the data that has been described, there is an increase in the number of Diabetes Mellitus sufferers which has a major influence on the increase in complications in Diabetes Mellitus patients.

There are generally more cases of diabetes mellitus occurs more often in developing countries than developed countries (Infodatin, 2018). According to Basic Health Research (2023) in Southeast Asia region, there are 82 million cases of diabetes mellitus in 2022 and is expected to increase to 151 million in 2045. The increase in cases continues to occur every year in the Southeast Asia region ranks third in the world. In Indonesia This case of diabetes mellitus also occurs frequently, this led to Indonesia being occupied ranked 6th out of 10 major countries with The highest number of diabetes mellitus cases in Asia (Basic Health Research, 2023)

One complication that can cause problems is wounds on the feet that cause infection or diabetic ulcers. The method of using Modern Dressing with Moist Wound Healing in the diabetic ulcer treatment process can speed up the wound healing process. Moist Wound Healing is a method that aims to keep the wound area always moist by using a moist dressing, or moist by maintaining the isolation of the wound environment made from occlusive and semi-occlusive materials. Moist Wound Healing supports the wound healing process so that natural tissue growth occurs which is moist and can expand if the amount of esudate is excessive, and prevents bacterial contamination from the outside.

Moist Wound Healing can speed up the wound healing process by 45% and can suppress the problem of inflammation and the development of scar tissue (Irwan et al., 2022). Based on research from Angriani et al., 2019 which states that healing of diabetic ulcers is effective, wound care is carried out using the moist wound healing method (Angriani et al., 2019). In line with research conducted by Oset et al., 2018 which states that the moist wound healing method tends to speed up the healing process of diabetic ulcers (Ose et al., 2018).

Based on the results of researchers' observation, Sering Road has experience in treating wounds using moist wound healing dressing techniques and has made many developments in the wound healing period, and many diabetes mellitus patients have had successful treatment. In interviews conducted by researchers with respondents who were undergoing wound care, they said that Sering Road more as a place to treat diabetes wounds, because the treatment was more modern and facilitated well. Based on this description, researchers are interested in conducting research on the effectiveness of modern wound care using the Moist Wound Healing Bandage Technique in Healing Diabetic Ulcers at the Sering Road.

METHODS

The method used in this research is quantitative research with a quasi-experimental design with a cross-sectional method approach. This design was used because researchers wanted to see the effectiveness of modern wound care using the Moist Wound Healing Bandage Technique in Healing Diabetic Ulcers at the Jl. Sering No. 20, Sidorejo, Kecamatan Medan Tembung, Kota Medan, Sumatera Utara 20222. The research location was carried out at the Sering Road. This research was carried out from January 01, 2024 to March 01, 2024. The population in this study were all patients who underwent wound care at Sering Road. The sample in this study amounted to 30 people. The sampling technique in this research is nonprobability sampling with a sampling method, namely accidental sampling. The accidental sampling method is an accidental sampling method by taking cases or respondents who

happen to be present or available in a place according to the research context, so that researchers can take samples from anyone they meet without prior planning.

RESULT AND DISCUSSION

Description of the characteristics of respondents based on research results at the Sering Road. The research results showed that the distribution of respondents' characteristics based on age showed that the majority of respondents were aged 56-65 years (late elderly) as many as 9 respondents (30.0%), this research is in line with the results of research conducted by (Primadani & Safitri, 2021) stating that age Respondents who healed diabetic foot wounds using the moist wound healing method were aged 40-60 years.

The distribution of respondents based on their latest level of education shows that the majority of respondents have a high school education, 13 respondents (43.3%). The distribution of respondents based on occupation shows that 9 respondents (30.0%) work as Housewives (IRT). The distribution of respondents based on length of wound care showed that 29 respondents (96.7%) had a length of wound care < 5 years. Based on the type of necrotic tissue, the majority of respondents had no necrotic tissue type, 24 respondents (80.0%).

This research is in line with research conducted by (Harmiady et al., 2020) which stated that the type of necrotic tissue in respondent 1 started with a score of 4, namely sticky, black false scar tissue, to a score of 2 in the seventh treatment, namely white/grey, slough. easy to remove, whereas for respondent 2 the initial score was 5, sticky with firm borders, hard and had black aschar, in the seventh treatment the score was 2, namely white/grey, slough was easy to remove. Based on the amount of necrotic tissue, the majority of respondents had no necrotic tissue, 23 respondents (76.7%).

Based on the amount of exudate, the majority of respondents had a small amount of exudate: the wound surface was moist, the exudate soaked < 25% of the dressing, 11 respondents (36.7%). This research is in line with research conducted by (Harmiady et al., 2020) which stated that the amount of exudate was large, it became bloody with little exudate and the wound looked damp at the seventh meeting.

Based on the skin color around the wound, the majority of respondents had pink skin color or normal skin color for each part of the wound, 18 respondents (60.0%). This research is in line with research conducted by (Harmiady et al., 2020) which stated that the skin color around the wound for the two respondents at the first meeting was a score of 5 and at the seventh meeting a score of 3, namely a change from hyperpigmentation to hypopigmentation or grayish.

Based on peripheral tissue induration, the majority of respondents had no induration, 30 respondents (100.0%). This research is in line with research conducted by (Harmiady et al., 2020) which stated that the hardening of peripheral tissue saw a decrease of 2 levels in each respondent, namely respondent 1 from a score of 5 to 3 and respondent 2 from a score of 4 to 2 in the seventh treatment where for score 5, namely hardening > 4cm in the area around the wound, score 4 hardening 2-4cm > 50% around the wound, score 3 < 50% around the wound and score 2 hardening < 2 cm around the wound.

Based on granulation tissue, the majority of respondents had bright granulation tissue, red like flesh; 75% to 100% of wounds were filled by 16 respondents (53.3%). This research is in line with research conducted by (Harmiady et al., 2020) which stated that the granulation tissue of the two respondents experienced a decrease from the first treatment with a score of 4 to 2 on the seventh day of treatment. It was seen that the granulation tissue from the first treatment was pink or dull or blackish, filled < 25% of the wound, while in the seventh treatment the bright red granulation color showed excess tissue growth which filled 75 to 100% of the wound.

Based on epithelialization, the majority of respondents had epithelialization < 25% with 11 respondents (36.7%). This research is in line with research conducted by (Harmiady et al., 2020) which stated that epithelialization from a score of 5 to 4 from the two respondents who were given treatment where < 25% of the wound was closed became 25% to < 50% of the wound was closed in the seventh treatment. The results of this study are inline with the results of research conducted by (Ose et al., 2018) regarding the effectiveness of wound care with wet-dry and moist wound healing techniques in healing diabetic ulcers which was carried out using the hydrocolloid dressing method. It was found that with the moist wound technique The healing process is faster than using wet-dry.

9 CONCLUSION

Based on the results of research and discussion, it can be concluded that:

1. Based on the research results, the characteristics of respondents based on age show that the majority of respondents are 56-65 years old (late elderly) as many as 9 respondents (30.0%), based on gender it shows that the majority of respondents are male, 17 respondents (56.7%), based on the latest level of education, it shows that the majority of respondents have a high school education as many as 13 respondents (43.3%), based on occupation shows that the respondents work as Housewives (IRT) as many as 9 respondents (30.0%), based on the length of wound care shows that the respondents have Wound care < 5 years was 29 respondents (96.7%).
2. Based on research results, there is the effectiveness of modern wound care using moist wound healing dressing techniques in healing diabetic ulcers at the Sering Road

REFERENCES

- Angriani, S., Hariani, H., & Dwianti, U. (2019). The effectivity of modern dressing wound care with moist wound healing method in diabetic ulcer at wound care clinic of etn centre Medan. *Jurnal Media Kesehatan*, 10(01), 19–24.
- Basic Health Research. 2023
- Detty, A. U., Fitriyani, N., Prasetya, T., Florentina, B. (2020). Karakteristik Ulkus Diabetikum Pada Penderita Diabetes Melitus The Characteristics of Diabetic Ulcer in Patients with Diabetes Mellitus. *Juni*, 11(1), 258–264.
<https://doi.org/10.35816/jiskh.v10i2.261>
- Fitria, E., Nur, A., Marissa, N., Ramadhan, N. (2017). Karakteristik Ulkus Diabetikum pada Penderita Diabetes Mellitus di RSUD dr. Zainal Abidin dan RSUD Meuraxa Banda Aceh. *Buletin Penelitian Kesehatan*, 45(3), 153–160. <https://doi.org/10.22435/bpk.v45i3.6818.153-160>.
- Harmiady, R., Ahmad, A. K., Putri, K. E., & Nasrullah, N. (2020). Efektivitas Metode Perawatan Luka “Moisture Balance” Terhadap Penyembuhan Luka Pada Pasien Ulkus Diabetikum Di Klinik Perawatan Luka Isam Cahaya Holistic Care Kota Medan. *Media Keperawatan: Politeknik Kesehatan Medan*, 11(2), 45. <https://doi.org/10.32382/jmk.v11i2.1942>.
- Irwan, M., Indrawati, Maryati, Risnah, & Arafah, S. (2022). Efektivitas Perawatan Luka Modern dan Konvensional terhadap Proses Penyembuhan Luka Diabetik. *Jurnal Ilmiah Mappadising*, 4(1), 237–245. <https://doi.org/10.54339/mappadising.v4i1.291>
- Kemendes RI. (2018). Hasil Riset Kesehatan Dasar Tahun 2018. *Kemendagri Kesehatan RI*, 53(9), 1689–1699.
- Kemendagri Kesehatan RI. (2020). Infodatin tetap produktif, cegah, dan atasi Diabetes Melitus 2020. In *Pusat Data dan Informasi Kemendagri Kesehatan RI* (pp. 1–10).
- Mutiudin, A. I. (2019). Efektivitas Proses Penyembuhan Luka dengan Penggunaan Modern Wound Dressing pada Pasien Ulkus Diabetik: A Sistematis Review. *Jurnal Keperawatan & Kebidanan*, 3(2), 12–21.
http://jmkubk.id/index.php/jmk_kb/article/view/84
- Rassi, Y.E.T., Efendy, M. A. (2023). Pola Diet Dengan Proses Penyembuhan Luka Diabetik. *Journal of Health Science Community*, 3(3), 113–117.
- Rusdi, M. S. (2020). Hipoglikemia Pada Pasien Diabetes Melitus. *Journal Syifa Sciences and Clinical Research*, 2(September), 83–90.
- Siregar, H. K., Batubara, K., Empraninta, H. E., & Pabanne, F. U. (2021). Relationship between Knowledge and Attitude about Prevention of Diabetes Mellitus Complications in Diabetes Mellitus Patients. *South East Asia Nursing Research*, 3(4), 153. <https://doi.org/10.26714/seanr.3.4.2021.153-155>

Application Of Modern Wound Care Using Moist Wound Healing Techniques In Diabetes Mellitus Patients

ORIGINALITY REPORT

11%

SIMILARITY INDEX

10%

INTERNET SOURCES

4%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	garuda.kemdikbud.go.id Internet Source	2%
2	asploro.com Internet Source	1%
3	Qatrinnada Maulidya Akmal, Hanina, Humaryanto, Lipinwati, Rita Halim. "ISOLATION AND IDENTIFICATION OF BACTERIA CAUSING SECONDARY WOUND INFECTIONS IN DIABETES MELLITUS PATIENTS IN JAMBI CITY", Jurnal Kedokteran dan Kesehatan : Publikasi Ilmiah Fakultas Kedokteran Universitas Sriwijaya, 2024 Publication	1%
4	www.midwifery.iocspublisher.org Internet Source	1%
5	web.emc.id Internet Source	1%
6	jurnal.akperrscikini.ac.id Internet Source	1%

7	jurnal.uimedan.ac.id Internet Source	1 %
8	Eko Naning Sofyanita, Anisa Rizky Yuniarti. "Effectiveness of white guava leaves (<i>Psidium guajava</i> var. <i>Pyrifera</i> L.) In repair proximal tubule damage and glomerulus diameter in hyperglycemic mice", ITEGAM- Journal of Engineering and Technology for Industrial Applications (ITEGAM-JETIA), 2023 Publication	<1 %
9	journal.uwks.ac.id Internet Source	<1 %
10	jurnal.unissula.ac.id Internet Source	<1 %
11	repository.politeknikyakpermas.ac.id Internet Source	<1 %
12	journal.poltekkes-mks.ac.id Internet Source	<1 %
13	j-innovative.org Internet Source	<1 %
14	Irwansyah Irwansyah, Ilcham Syarief Kasim. "Identifikasi Keterkaitan Lifestyle Dengan Risiko Diabetes Melitus", Jurnal Ilmiah Kesehatan Sandi Husada, 2021 Publication	<1 %

ejournal.nusantaraglobal.ac.id

15	Internet Source	<1 %
16	masjidjaminurulihsan.blogspot.com Internet Source	<1 %
17	repository.stikeshangtuah-sby.ac.id Internet Source	<1 %
18	www.journal.stikespemkabjombang.ac.id Internet Source	<1 %
19	akper-sandikarsa.e-journal.id Internet Source	<1 %
20	ijisrt.com Internet Source	<1 %
21	www.dovepress.com Internet Source	<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Application Of Modern Wound Care Using Moist Wound Healing Techniques In Diabetes Mellitus Patients

GRADEMARK REPORT

FINAL GRADE

GENERAL COMMENTS

/0

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
